

Precise measurement of HFS of positronium


A. Ishida*, G. Akimoto*, K. Kato*, T. Suehara*, T. Namba*, S. Asai*, T. Kobayashi*,
H. Saito[†], M. Yoshida**, K. Tanaka**, A. Yamamoto**, I. Ogawa[‡], S. Kobayashi[‡] and T. Idehara[‡]

University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan
^{*}Department of Physics and ICEPP, [†]Institute of Physics, University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8902, Japan
^{**}High Energy Accelerator Research Organization (KEK), 1-1 Oho, Tsukuba, Ibaraki, 305-0801, Japan
[‡]FIR Center, University of Fukui, 3-9-1 Bunkyo, Fukui, 910-8507, Japan

Positronium and its hyperfine structure (HFS)

Positronium (Ps)

The bound state of an electron (e^-) and a positron (e^+)
 orthopositronium (o-Ps) $\cdots 1^3S_1$ mostly 3γ decay
 parapositronium (p-Ps) $\cdots 1^1S_0$ mostly 2γ decay



Hyperfine structure (HFS)

- The energy splitting between o-Ps and p-Ps
- The value of HFS


Experimental average

203.388 65(67) GHz (3.3 ppm)
 PRA 27, 262 (1983)
 PRA 30, 1331 (1984)

Theory

203.391 69(41) GHz (2.0 ppm)
 PRL 85, 5094 (2000)

- The measured values are **consistent with each other** and **lower than the theoretical calculation**.



15 ppm (3.9 σ) discrepancy


Measurement using the Zeeman effect

How to measure the HFS?

Induce the transition

$\rightarrow 2\gamma$ decay rate increases.

- In a static magnetic field, energy levels of o-Ps split between $m_z=0$ and $m_z=\pm 1$ states. (**Zeeman Effect**)
- At about **9 kG**, Δ_{mix} is about **3 GHz (microwave)**.
- The HFS value is **calculated from Δ_{mix}** . (**indirect measurement**)
- What about direct measurement?
 \rightarrow See T. Suehara's poster.



Common systematic uncertainties in the previous experiments

1. Underestimation of material effects

- Unthermalized o-Ps can have a significant effect (especially at low material density). \leftarrow o-Ps lifetime puzzle (1990's)

2. Non-uniformity of the magnetic field

- It's quite difficult to get ppm level uniform field in a large Ps creation volume

Experimental setup

To **reduce these systematic uncertainties**, we use the following **new methods**.

Large bore superconducting magnet

- Operated in **Persistent Current mode** (stable).
- **70 ppm magnetic field uniformity** without any compensations.

Waveguide

2.853 GHz
500 W CW RF
 (GaN Amplifier)

80 cm

RF Cavity
 $f_0 = 2.853 \text{ GHz}$
 $Q = 14700 \pm 50$
 TM₁₁₀ mode
 Filled with 1.5 atm gas (90% N₂ + 10% iso-C₄H₁₀)

208 mm

High performance gamma-ray detectors

LaBr₃ (Ce) scintillators (x 6)
 1.5" in diameter & 2.0" long

High energy and timing resolutions, short decay constant


Time information

- Plastic scintillator is used to **tag emitted β^+** .
- Get the time information between o-Ps creation ($t = 0$) and decay.

(1) We can measure the thermalization.

(2) Prompt suppression

0.2 mm thick, 15 mm x 15 mm Plastic Scintillator



Inside of the RF cavity


²²Na source (700 kBq)

Current status

We are presently taking more data....


Preliminary plots

ENERGY SPECTRA




2γ decay rate increases because of the transition between o-Ps' $m_z=0$ and $m_z=\pm 1$ states.

TIMING SPECTRA



RESONANCE CURVE



Converted HFS value (from an **only 2 weeks run**) is **203.399**

± 0.005 (23 ppm, stat.)


± 0.029 (140 ppm, sys.) GHz (Preliminary)

(consistent with the previous experiments)
 The systematic error mainly comes from the non-uniformity of the magnetic field.

Our goal

O (1) ppm accuracy in a year

1. Develop compensation coils
 \rightarrow **Get O(1) ppm B - uniformity**



2. Precisely measure the thermalization function.

3. **Derive the HFS value at O (1) ppm accuracy.**

\rightarrow **Solve or Confirm the discrepancy between the experimental values and the theoretical value.**